

Comparison of the Effects of Marine Omega-3 Fatty Acids and Ibuprofen on Primary DysmenorrheaKooshki A^{1*}, Tofighiyan T², Rakhshani M H².¹. Traditional and Complementary Medicine Research Center, Sabzevar University of Medical Sciences, Sabzevar, Iran². Department of nursing, Sabzevar University of Medical Sciences, Sabzevar, IranE-mail: akooshki.nutr@yahoo.com, Tell: +98-0910-2911161

Abstract: Dysmenorrhea is common gynecological problem among women. This study was done to examine the effects omega-3 fatty acid compared to ibuprofen on dysmenorrhea. A double- blind crossover study was conducted among 18-22 years old female student. Women assigned to group 1 received 2 omega-3 capsules daily in the morning and night after food for 2 month and group 2 receive 400 mg ibuprofen very 8 hours in three days. It was a one month washout period. Then each group received drug other group. Data were gathered the start, after 8 and 20 weeks by using the Cox Menstrual Symptom Scale questionnaire. Data were analyzed using spss 16.0 via Paired t- test and Repeated Measurement. The mean age of students was 20.35 ± 1.03 years and their mean BMI was 21.03 ± 2.40 kg /m². Result showed a marked reduction in pain intensity after treatment with omega-3 ($P < 0.05$). Also our results indicate that in the first day ($P = 0.28$), the second day ($P = 0.30$) and the third day ($P = 0.67$) of menstruation, the effect of omega-3 and ibuprofen was similar. The results indicate that dietary supplementation with omega-3 fatty acid can reduced noticeable pain as ibuprofen among young women with primary dysmenorrhea.

[Kooshki A, Tofighiyan T, Rakhshani M H. **Comparison of the Effects of Marine Omega-3 Fatty Acids and Ibuprofen on primary dysmenorrhea.** *Life Sci J* 2013;10(4s):1-3] (ISSN:1097-8135) <http://www.lifesciencesite.com>. 1

Key word: Dysmenorrhea, Omega-3, ibuprofen**1. Introduction**

Dysmenorrhea is common gynecological problem among women which is divided into two categories, primary and secondary. Primary dysmenorrhea is painful menstruation in the absence of pelvic disease. The pain of uterine origin is associated with symptoms such as nausea, vomiting, abdominal distension, diarrhea, headache and rarely even myocardial infarction (1,2).

Prevalence of primary dysmenorrhea in various communities should have between 50 to 90% (3,4). The prevalence of dysmenorrhea have been reported 74-86% in Iran (4)

The use of non- steroidal anti- inflammatory drugs (NSAIDS) can reduce pain of primary dysmenorrhea in women with primary dysmenorrhea(5). There fore, adverse side effects may occur after long- term use of these drugs. These adverse effects are impaired kidney function and gastrointestinal bleeding which could limit their use in primary dysmenorrhea(6,7).

Today, various studies dietary supplements have been effective for the treatment of dysmenorrhea. These supplements are omega-3 fatty acids, magnesium, zinc, vitamin B and vitamin E (8). Research has shown omega-3 two months can receive a blood concentration of this substance to induce anti-inflammatory and analgesic effects provided (9,10,11,12). Most studies have shown that omega-3 fatty acids are well tolerated by most people and

rarely in the high dose (more than 6 grams per day) are adverse side effects (13). Omega-3 fatty acids possess anti- inflammatory effects due to produce vasodilator eicosanoids such as PGE3 and PGF3 and inhibit of production vasoconstrictor prostaglandin PGE2, PGF2 from omega-6 which is the important cause primary dysmenorrhea(12,14).

The aim of the study was to evaluate the effects of marine omega-3 fatty acids compared to ibuprofen on primary dysmenorrhea.

2. Materials and methods

A double- blind crossover study was conducted among 18-22 years old female student at Sabzevar university of Medical science. The students had primary menstrual pain without significant pathology and with regular menstrual cycles. The research proposal was approved by the Ethics committee of Sabzevar university of Medical science and all students signed a consent form.

The sample size estimation for each group was 24 at a power of $1 - \beta$ of 90% and $\alpha = 0.05\%$ and C.V 10% and $t(0.025, 26) = 2.056$ and $t(0.1, 26) = 1.315$ based on below formula:

$$n \geq \{t(\alpha/2, 2n-2) + t(\beta, 2n-2)\}^2 \{C.V/10\}^2 \quad (15).$$

After a primary examination, women who had inclusion criteria were participated in the present study. All student divide to two groups based on table of random numbers.

Women in group 1 received 2 omega-3 capsule (contain of 180 mg of EPA and 120 mg of DHA) daily in the morning and night after food for 2 month and group 2 receive 400mg ibuprofen a very 8 hours in three days.(One day before and two days after menstrual period three times in day). It was a one month washout period without any medication. Then each group received drug other group.

After the time, Data were gathered before the start, after 8 and 20 weeks by using the Cox Menstrual Symptom Scale questionnaire (16).In addition, the dietary intakes of subjects were assessed a 2-days dietary recall at baseline and the end of weeks 8 and 20. Subject diets were analyzed by Nutrition-IV software. Data were analyzed using SPSS version 16.0 via Paired t- test and Repeated Measurement.

3. Results

of the 50 female student who enrolled, 3 were lost to follow- up. The mean age of students and mean age

of first menorrh and dysmenorrhea was shown in table 1. 56% of all students in first day, 18% in second day and 16% in third day had pain.

Table 1: characteristics of study subjects

Mean \pm SD	
Age (year)	20.35 \pm 1.03
Age first menorrh (years)	13.47 \pm 2.29
Age of dysmenorrh (years)	12.26 \pm 5.67
BMI (kg/m ²)	21.03 \pm 2.40

AS shown in table 2, the mean pain severity score was significantly reduced after omega-3 supplementation among female students (P=0.002). Also, our results indicate that in the first day (P=0.28), the second day(P=0.30) and the third day(P=0.67) of menstruation, the effect of omega-3 and ibuprofen was similar. Anthropometric and dietary factors were not significantly different at baseline and the end of weeks 8.

Table 2: Mean \pm SD of pain after intake of omega and Ibuprofen

Omega group		Ibuprofen group	
before	after	before	after
5.89 \pm 3.04	7.67 \pm 1.49	5.8 \pm 2.7	7.7 \pm 1.52
P-value: 0.002		P-value: 0.0001	

4. Discussion

In the present study, omega-3 fatty acid supplementation was reduced dysmenorrheal in female students but this reduction pain was not significant comparison ibuprofen. This results of study is consistent with findings Dowlatiyan(1). Dowlatiyan et al showed that dietary supplementation with omega-3 fatty acids significantly decreased primary dysmenorrhea. However the present study design omega-3 was compared to ibuprofen. Rahbar et al showed that dietary supplementation with omega-3 fatty acids significantly decreased primary dysmenorrhea. However the present study design was higher dosage of omega-3 (2 versus 1capsule) and included a washout period and compared to ibuprofen (8).

In another study, Zamani et al evaluated the effect of dietary supplementation with omega-3 fatty acids on primary dysmenorrhea on 84 adolescents(9). The first group received 1 gram omega-3 supplement daily for 2 months and second group received placebo in the same time. Results of their study showed that omega-3 fatty acids supplement significantly reduced intensity and duration of pain in adolescent girls.

Moghadamnia et al showed that omega-3 fatty acids significantly decrease abdominal and low-back

pain and need for NSAIDs drug in adolescent girls with primary dysmenorrheal(17).

Safari et al assessed the effect of dietary supplementation with vitamin E on primary dysmenorrhea on 86 female students based to randomized design (18). The first group received 500 IU vitamin E supplement daily for 2days before and 1 day after menarche and second group received 250 mg of mefenamic acid every 8 hours in the same time. Results of their study showed that reduction of pain was not significant between two group. In other words, vitamin E and mefenamic acid reduced dysmenorrhea to a same value. The finding reported by Safari are similar to our results quality impact of dietary supplement with NSAIDs.

Finally, Detuch et al assessed the influence intake of fish and sea foods and menstrual pain on 220 women 20-45 years old(19). He found that low consumption of fish and sea foods correlated with primary dysmenorrheal. The other words, women had highest level of intake of fish in their diet, they not dysmenorrhea.

The results of the present study indicate that dietary supplementation with omega-3 fatty acid can reduced noticeable pain as ibuprofen among young women with primary dysmenorrhea. Nevertheless, the use of omega-3 fatty acids could potentially offer

for the safe and effective treatment of women with dysmenorrheal compared to NSAIDS.

Acknowledgement:

The present study was supported by Sabzevar university of Medical Sciences(No:11609_ xxx). The authors also gratefully acknowledge the cooperation of participating patients, without whom this research would not have been possible.

Corresponding Author:

Akram Kooshki,
Traditional and Complementary Medicine Research
Center, Sabzevar University of Medical Sciences,
Sabzevar, Iran
E-mail:akooshki.nutr@yahoo.com

References

1. Dowlatiyan M, Jafari H, Valaei N, Afrakhteh M, Taleban FA, Gachkar L. Evaluation of the effects of fish oil on primary dysmenorrhea. J Zanzan Uni Med Scien 2004;12(47):7-13
2. Proctor ML, Farquhar CM. Dysmenorrhea. Clin Evid 2006;15:2429-48
3. Berek JS, Adashi E, Hillard PA. Novaks Gynecology. 13th ed. USA: Williams & Willkins, Mosby Co:2002.421-39
4. Khooshechin T. The effect of vitamin E on primary dysmenorrheal. Modares J Med Scien 1998; 1(1):13-15
5. Own PR. Prostaglandin synthetase inhibitors in the treatment of primary dysmenorrheal. Am J Obstet Gynecol 1984;148(1):96-103
6. Drevon CA. Marine oils and their effects. Nutr Rev 1992;50(4 suppl 2):38-45
7. Smith WL, Langenbach R. Why there are two cyclooxygenase isozymes. J Clin Invest 2001;107(12):1491-5
8. Rahbar N, Asgharzadeh N, Ghorbani R. Effect of omega-3 fatty acids on intensity of primary dysmenorrhea. Int J Gynaecol Obstet. 2012 Jan 17. [Epub ahead of print]
9. Zamani M, Arab M, Nasrollahi Sh, Kashani Kh. The effects of fish oil on primary dysmenorrhea. J Gorgan Uni Med Scien 2005;7(1):39-42
10. Proctor ML, Murphy PA. Herbal and dietary therapies for primary and secondary dysmenorrheal. Cochrane Database Syst Rev. 2001;3:2124
11. Fjerbaek A, Knudsen UB. Endometriosis, dysmenorrhea and diet- What is the evidence? Eur J Obstet Gynecol Reprod Biol 2007;132(2):140-147
12. Saldeen P, Saldeen T. Women and omega fatty Acids. Obstet Gynecol Surv 2004;59(10):722-730.
13. Friedman A, Moe SH. Review of effects of omega-3 supplementation in dialysis patients. Clin Am Soc Nephrol. 2006; 1:182-192.
14. Harel Z, Biro FM, Kottenhahn RK, Rosenthal SI. Supplementation with omega-3 polyunsaturated fatty acids in the management of dysmenorrhea in adolescences. Am J Obstet Gynecol 1996;174(4):1335-8
15. Chow SC and Liu JP. Design and analysis of clinical trials, concepts and methodologies. Second edition, Newjersey, John Wiley and sons, 2004, Chapter 11, section 7.
16. Cox DJ, Meyer RG. Behavioral treatment parameters with primary dysmenorrheal. J Behav Med 1978;1(3):297-310
17. Moghadamnia AA, Mirhosseini N, Abadi MH, Omranirad A, Omiavar S. Effect of Clupeonella grimmii fish oil on dysmenorrheal. East Mediterr Health J 2010;16(4):408-413
18. Safari A, Shah Rezaei Gh, Damavandi A. Comparison of the effects of Mefnamic acid and vitamin E on primary dysmenorrhea. J Army uni Med Scien 2006; 4(1):742-738
19. Detuch B. Painful menstruation and low intake of w-3 fatty Acids. Ugeskr laeger 1996;158(29):4195-4198

1/12/2013